Author Index

Abbiati, A., see Guiso, G. 363
Akesson, B.
— , Forslind, K. and Wollheim, F.
Analysis of homogentisic acid in body fluids by high-performance liquid chromatography 233
Ang, S.B., see Lee, E.J.D. 300
Anzion, R.B.M., see Jongeneelen, F.J. 227
Assenza, S.P., see Koenigbauer, M.J. 161
Baba, S., see Furuta, T. 1
Bayne, W.F., see Constanzer, M.L. 277
— , see Stubbs, R.J. 171
Belleville, F., see Brossat, B. 141
Birckel, P., see Jehl, F. 109
Botha, H., see Swart, K.J. 315
Brimmell, P.A.
— and Sams, D.J.
Rapid and simple assay for the measurement of methotrexate in serum, urine and red blood cells by reversed-phase high-performance liquid chromatography 320
Brossat, B.
— , Straczek, J., Heulin, M.-H., Herbeuval, X., Belleville, F., Nabet, P. and Lokiec, F.
Liquid chromatographic determination of cyclosporin A in serum with use of a solid-phase extraction. Comparison between high-performance liquid chromatography and radioimmunoassay levels in clinical investigations 141
Caccia, S., see Guiso, G. 363
Chakrabarti, S., see Colin, P. 151
Christians, U.
— , Zimmer, K.-O., Wonigeit, K. and Sewing, K.-Fr.
Measurement of cyclosporin A and of four metabolites in whole blood by high-performance liquid chromatography 121
Colacchio, T.A., see Nierenberg, D.W. 79
Colin, P.
— , Sirois, G. and Chakrabarti, S.
Rapid high-performance liquid chromatographic assay of acetaminophen in serum and tissue homogenates 151
Collins, A.J., see Notarianni, L.J. 305
Constanzer, M.L.
— and Bayne, W.F.
Determination of picogram levels of (*E*)-3-(9-chloro-6,11-dihydro-5H-pyrrolo(2,1-B)(3)benzazepin-11-ylidene)-N,N-dimethyl-1-propanamine (*Z*)-2-butenedioate (1:1) in plasma using capillary gas chromatography with nitrogen-selective detection 277
Cukrová, V.
— and Hrkal, Z.
Purification and characterization of granulocyte-macrophage colony stimulating factor from human placenta 242
Curtis, M.A., see Koenigbauer, M.J. 161
Czejka, M.J.
— and Koch, H.P.
Determination of thalidomide and its major metabolites by high-performance liquid chromatography 181
Danhauser, L., see Gandhi, V. 293
Dubois, J.P., see Sioufi, A. 101
Eibel, G., see Scharpf, F. 91
Elchisak, M.A., see Scott, M.C. 17
Entwistle, L.A., see Stubbs, R.J. 171
Er, S.S., see McGarrity, M.J. 53
Forgione, A., see Pedrazzini, S. 338
Forslind, K., see Akesson, B. 233
Fracasso, C., see Guiso, G. 363
Furuta, T.
— , Kasuya, Y., Shibasaki, H. and Baba, S.
Determination of stable isotopically substituted histidine in human plasma by gas chromatography—mass spectrometry 1
Gandhi, V.
— , Danhauser, L. and Plunkett, W.
Separation of 1-β-D-arabinofuranosylcytosine 5′-triphosphate and 9-β-D-arabinofuranosyl-2-fluoroadenine 5′-triphosphate in human leukemia cells by high-performance liquid chromatography 293
Garvie, C.T.
— , Straub, K.M. and Lynn, R.K.
Quantitative liquid chromatographic determination of disulfide-containing peptide analogues of vasopressin with dual Hg/Au electrochemical detection 43
Gherezghiher, T.
— , Koss, M.C., Nordquist, R.E. and Wilkinson, C.P.
Rapid and sensitive method for measurement of hyaluronic acid and isomeric chondroitin sulfates using high-performance liquid chromatography 9

Granneman, G.R.
— and Sennello, L.T.
High-performance liquid chromatographic procedures for the determination of difloxacin and its metabolites in biological matrices 199
Griffin, R.L., see Minkler, P.E. 33
Guiso, G.
— , Fracasso, C., Caccia, S. and Abbiati, A.
Determination of ranitidine in rat plasma and brain by high-performance liquid chromatography 363
Gyllenhaal, O.
Derivatization of 2-amino alcohols with phosgene in aqueous media: limitations of the reaction selectivity as found in the presence of O-glucuronides of alprenolol in urine 270
Hada, T., see Yamamoto, T. 342
Haginaka, J.
— and Wakai, J.
Liquid chromatographic determination of amoxicillin and its metabolites in human urine by postcolumn degradation with sodium hypochlorite 219
Hammes, W.
— and Weyhenmeyer, R.
Quantitative determination of vinpocetine in human plasma by capillary gas chromatography—mass spectrometry 264
Hayakawa, K.
— and Oizumi, J.
Determination of free biotin in plasma by liquid chromatography with fluorimetric detection 247
Henderson, P.Th., see Jongeneelen, F.J. 227
Herbeuval, X., see Brossat, B. 141
Heulin, M.-H., see Brossat, B. 141
Heykants, J., see Woestenborghs, R. 332
Higashino, K., see Yamamoto, T. 342
Hoppel, C.L., see Minkler, P.E. 33
Hrkal, Z., see Cukrová, V. 242
Hsia, J.C., see McGarrity, M.J. 53
Ingalls, S.T., see Minkler, P.E., 33
Iwamura, S.
— and Tsukamoto, K.
Highly sensitive determination of 2-hydroxy-N-{3-[*m*-(1-piperidinylmethyl)-phenoxy]propyl}acetamide (desacetyl TZU-0460) in rat plasma by high-performance liquid chromatography 370
Jaeger, H., see Nieder, M. 207
Jehl, F.
— , Birckel, P. and Monteil, H.
Hospital routine analysis of penicillins, third-generation cephalosporins and aztreonam by conventional and high-speed high-performance liquid chromatography 109
Jongeneelen, F.J.
— , Anzion, R.B.M. and Henderson, P.Th.
Determination of hydroxylated metabolites of polycyclic aromatic hydrocarbons in urine 227
Kasuya, Y., see Furuta, T. 1
Katae, H., see Kojima, K. 326
Kleinjans, J.C.S., see Verhagen, H. 282
Koch, H.P., see Czejka, M.J. 181
Koenigbauer, M.J.
— , Assenza, S.P., Willoughby, R.C. and Curtis, M.A.
Trace analysis of diazepam using microbore high-performance liquid chromatography and on-line preconcentration 161
Kojima, K.
— , Yamamoto, K., Nakanishi, Y. and Katae, H.
Determination of 22,23-dihydroavermectin B_{1a} in dog plasma using solid-phase extraction and high-performance liquid chromatography 326
Komoda, T., see Koyama, I. 65
Koss, M.C., see Gherezghiher, T. 9
Koyama, I.
— , Miura, M., Matsuzaki, H., Sakagishi, Y. and Komoda, T.
Sugar-chain heterogeneity of human alkaline phosphatases: differences between normal and tumour-associated isozymes 65
Kraus, R.
— , Wuhte, J. and Rüfer, R.
Method for determination of different phospholipids by circular high-performance thin-layer chromatography 257
Lam, G.
— , Williams, R.M. and Whitney, C.C.
Electrochemical determination of oxymorphone in rat plasma by ion-pair reversed-phase high-performance liquid chromatography 309
Laufen, H., see Scharpf, F. 91
Lee, E.J.D.
— and Ang, S.B.
Simple and sensitive high-performance

liquid chromatographic assay for 5-aminosalicylic acid and acetyl-5-aminosalicylic acid in serum 300
Lemonnier, F., see Therasse, J. 237
Lesne, M., see Wallemacq, P.E. 131
Lester, D.C., see Nierenberg, D.W. 79
Lindgren, K.
Determination of cefaclor and cephradine in serum by ion-pair reversed-phase chromatography 351
— Determination of cefadroxil in serum by high-performance liquid chromatography with cephradine as internal standard 347
Lokiec, F., see Brossat, B. 141
Lorreyne, W., see Woestenborghs, R. 332
Lynn, R.K., see Garvie, C.T. 43
McGarrity, M.J.
— , Er, S.S., Nightingale, K.A. and Hsia, J.C.
Isolation and partial characterization of pyridoxal 5′-phosphate hemoglobins by high-performance liquid chromatography as a quality-control method for hemoglobin-based blood substitutes 53
Matsuzaki, H., see Koyama, I. 65
Midskov, C.
High-performance liquid chromatographic assay of fenflumizole and its demethyl metabolites in biological samples 189
Minkler, P.E.
— , Ingalls, S.T., Griffin, R.L. and Hoppel, C.L.
Rapid high-performance liquid chromatography of 3-methylhistidine in human urine 33
Misawa, S., see Tanaka, E. 376
Miura, M., see Koyama, I. 65
Monteil, H., see Jehl, F. 109
Moriwaki, Y., see Yamamoto, T. 342
Müller, S., see Reiter, C. 251
Müller, T., see Reiter, C. 251
Muramatsu, I., see Tamaki, M. 287
Nabet, P., see Brossat, B. 141
Nakanishi, Y., see Kojima, K. 326
Ng, L.L., see Stubbs, R.J. 171
Nieder, M.
— and Jaeger, H.
Quantification of tamoxifen and N-desmethyltamoxifen in human plasma by high-performance liquid chromatography, photochemical reaction and fluorescence detection, and its application to biopharmaceutic investigations 207
Nierenberg, D.W.
— , Lester, D.C. and Colacchio, T.A.
Determination of tocopherol and tocopherol acetate concentrations in human feces using high-performance liquid chromatography 79
Nightingale, K.A., see McGarrity, M.J. 53
Nordquist, R.E., see Gherezghiher, T. 9
Notarianni, L.J.
— and Collins, A.J.
Method for the determination of acemetacin, a non-steroidal anti-inflammatory drug, in plasma by high-performance liquid chromatography 305
Nozaki, S., see Tamaki, M. 287
Oizumi, J., see Hayakawa, K. 247
Pedrazzini, S.
— , Zanoboni-Muciaccia, W. and Forgione, A.
Determination of the non-steroidal anti-inflammatory drug flunoxaprofen, *S*(+)-2-(4-fluorophenyl)-α-methyl-5-benzoxazoleacetic acid, in blood and urine by high-performance liquid chromatography 338
Plunkett, W., see Gandhi, V. 293
Pommier, F., see Sioufi, A. 101
Reiff, K.
High-performance liquid chromatographic method for the determination of carvedilol and its desmethyl metabolite in body fluids 355
Reiter, C.
— , Müller, S. and Müller, T.
Improved method for the determination of sulphate in human serum using ion chromatography 251
Rüfer, R., see Kraus, R. 257
Sakagishi, Y., see Koyama, I. 65
Sams, D.J., see Brimmell, P.A. 320
Scharpf, F.
— , Yeates, R.A., Laufen, H. and Eibel, G.
Gas chromatographic assay of glycerol mononitrates in biological samples 91
Scott, M.C.
— and Elchisak, M.A.
Direct measurement of dopamine O-sulfate in plasma and cerebrospinal fluid 17
Senello, L.T., see Granneman, G.R. 199
Sewing, K.-Fr., see Christians, U. 121
Shibasaki, H., see Furuta, T. 1
Sioufi, A.
— , Pommier, F. and Dubois, J.P.
Determination of the two dinitrate

metabolites of nitroglycerin in human plasma by capillary gas chromatography with electron-capture detection 101
Sirois, G., see Colin, P. 151
Straczek, J., see Brossat, B. 141
Straub, K.M., see Garvie, C.T. 43
Stubbs, R.J.
— , Ng, L.L., Entwistle, L.A. and Bayne, W.F.
Analysis of sulindac and metabolites in plasma and urine by high-performance liquid chromatography 171
Swart, K.J.
— and Botha, H.
Rapid method for the determination of the diuretic triamterene and its metabolites in plasma and urine by high-performance liquid chromatography 315
Takahashi, S., see Yamamoto, T. 342
Takimoto, M., see Tamaki, M. 287
Tamaki, M.
— , Takimoto, M., Nozaki, S. and Muramatsu, I.
Adsorption of cyclic peptides analogous to gramicidin S and gratisin onto octadecylsilica stationary phase and bacterial cells 287
Tanaka, E.
— and Misawa, S.
Simultaneous determination of serum trimethadione and its metabolite by gas chromatography 376
Therasse, J.
— and Lemonnier, F.
Determination of plasma lipoperoxides by high-performance liquid chromatography 237
Thijssen, H.H.W., see Verhagen, H. 282
Tsukamoto, K., see Iwamura, S. 370
Työppönen, J.T.
Rapid and sensitive determination of Dns-amino acids in plasma using high-speed octadecyl liquid chromatographic columns 25
Verhagen, H.
— , Thijssen, H.H.W. and Kleinjans, J.C.S.
Sensitive high-performance liquid chromatographic method for the routine determination of butylated hydroxyanisole in plasma 282
Wakai, J., see Haginaka, J. 219
Wallemacq, P.E.
— and Lesne, M.
New automated high-performance liquid chromatographic analysis of cyclosporin A and G in human serum 131
131
Weyhenmeyer, R., see Hammes, W. 264
Whitney, C.C., see Lam, G. 309
Wilkinson, C.P., see Gherezghiher, T. 9
Williams, R.M., see Lam, G. 309
Willoughby, R.C., see Koenigbauer, M.J. 161
Woestenborghs, R.
— , Lorreyne, W. and Heykants, J.
Determination of itraconazole in plasma and animal tissue by high-performance liquid chromatography 332
Wollheim, F., see Akesson, B. 233
Wonigeit, K., see Christians, U. 121
Wuthe, J., see Kraus, R. 257
Yamamoto, K., see Kojima, K. 326
Yamamoto, T.
— , Moriwaki, Y., Takahashi, S., Hada, T. and Higashino, K.
Rapid and simultaneous determination of pyrazinamide and its major metabolites in human plasma by high-performance liquid chromatography 342
Yeates, R.A., see Scharpf, F. 91
Zanoboni-Muciaccia, W., see Pedrazzini, S. 338
Zimmer, K.-O., see Christians, U. 121

Subject Index

Acemetacin
Method for the determination of acemetacin, a non-steroidal anti-inflammatory drug, in plasma by HPLC 305

Acetaminophen
Rapid HPCL assay of acetaminophen in serum and tissue homogenates 151

Acetyl-5-aminosalicylic acid
Simple and sensitive HPLC assay for 5-aminosalicylic acid and acetyl-5-aminosalicylic acid in serum 300

Alkaline phosphatases
Sugar-chain heterogeneity of human alkaline phosphatases: differences between normal and tumour-associated isozymes 65

Alprenolol
Derivatization of 2-amino alcohols with phosgene in aqueous media: limitations of the reaction selectivity as found in the presence of O-glucuronides of alprenolol in urine 270

Amino acids
Determination of stable isotopically substituted histidine in human plasma by GC–MS 1

Amino acids
Rapid and sensitive determination of Dns-amino acids in plasma using high-speed octadecyl LC columns 25

Amino acids, methylated
Rapid HPLC of 3-methylhistidine in human urine 33

2-Amino alcohols
Derivatization of 2-amino alcohols with phosgene in aqueous media: limitations of the reaction selectivity as found in the presence of O-glucuronides of alprenolol in urine 270

5-Aminosalicylic acid
Simple and sensitive HPLC assay for 5-aminosalicylic acid and acetyl-5-aminosalicylic acid in serum 300

Amoxicillin
LC determination of amoxicillin and its metabolites in human urine by post-column degradation with sodium hypochlorite 219

Amoxicilloic acid
LC determination of amoxicillin and its metabolites in human urine by post-column degradation with sodium hypochlorite 219

1-β-D-Arabinofuranosylcytosine 5′-triphosphate
Separation of 1-β-D-arabinofuranosylcytosine 5′-triphosphate and 9-β-D-arabinofuranosyl-2-fluoroadenine 5′-triphosphate in human leukemia cells by HPLC 293

9-β-D-Arabinofuranosyl-2-fluoroadenine 5′-triphosphate
Separation of 1-β-D-arabinofuranosylcytosine 5′-triphosphate and 9-β-D-arabinofuranosyl-2-fluoroadenine 5′-triphosphate in human leukemia cells by HPLC 293

Aztreonam
Hospital routine analysis of penicillins, third-generation cephalosporins and aztreonam by conventional and high-speed HPLC 109

Biotin
Determination of free biotin in plasma by LC with fluorimetric detection 247

tert.-Butyl-4-hydroxyanisole
Sensitive HPLC method for the routine determination of butylated hydroxyanisole in plasma 282

Carvedilol
HPLC method for the determination of carvedilol and its desmethyl metabolite in body fluids 355

Cefaclor
Determination of cefaclor and cephradine in serum by ion-pair reversed-phase chromatography 351

Cefadroxil
Determination of cefadroxil in serum by HPLC with cephradine as internal standard 347

Cephalosporins
Hospital routine analysis of penicillins, third-generation cephalosporins and aztreonam by conventional and high-speed HPLC 109

Cephradine
Determination of cefaclor and cephradine in serum by ion-pair reversed-phase chromatography 351

Cephradine
Determination of cefadroxil in serum by HPLC with cephradine as internal standard 347

(*E*)-3-(9-Chloro-6,11-dihydro-5H-pyrrolo(2,1-B)(3)benzazepin-11-ylidene)-N,N-dimethyl-1-propanamine (*Z*)-2-butenedioate (1:1)
Determination of picogram levels of

(*E*)-3-(9-chloro-6,11-dihydro-5H-pyrrolo(2,1-B)(3)benzazepin-11-ylidene)-N,N-dimethyl-1-propanamine (*Z*)-2-butenedioate (1:1) in plasma using capillary GC with nitrogen-selective detection 277

Chondroitin sulphates
Rapid and sensitive method for measurement of hyaluronic acid and isomeric chondroitin sulfates using HPLC 9

Colony stimulating factor
Purification and characterization of granulocyte-macrophage colony stimulating factor from human placenta 242

Cyclosporins
New automated HPLC analysis of cyclosporin A and G in human serum 131

Cyclosporins
LC determination of cyclosporin A in serum with use of a solid-phase extraction. Comparison between HPLC and radioimmunoassay levels in clinical investigations 141

Cyclosporins
Measurement of cyclosporin A and of four metabolites in whole blood by HPLC 121

Demethylfenflumizoles
HPLC assay of fenflumizole and its demethyl metabolites in biological samples 189

Desmethylcarvedilol
HPLC method for the determination of carvedilol and its desmethyl metabolite in body fluids 355

N-Desmethyldifloxacin
HPLC procedures for the determination of difloxacin and its metabolites in biological matrices 199

N-Desmethyltamoxifen
Quantitation of tamoxifen and N-desmethyltamoxifen in human plasma by HPLC, photochemical reaction and fluorescence detection, and its application to biopharmaceutic investigations 207

Diazepam
Trace analysis of diazepam in serum using microbore HPLC and on-line preconcentration 161

Difloxacin
HPLC procedures for the determination of difloxacin and its metabolites in biological matrices 199

Difloxacin N-oxide
HPLC procedures for the determination of difloxacin and its metabolites in biological matrices 199

22,23-Dihydroavermectin B_{1a}
Determination of 22,23-dihydroavermectin B_{1a} in dog plasma using solid-phase extraction and HPLC 326

Dimethadione
Simultaneous determination of serum trimethadione and its metabolite by GC 376

Dns-amino acids
Rapid and sensitive determination of Dns-amino acids in plasma using high-speed octadecyl LC columns 25

Dopamine O-sulfate
Direct measurement of dopamine O-sulfate in plasma and cerebrospinal fluid 17

Enzymes
Sugar-chain heterogeneity of human alkaline phosphatases: differences between normal and tumour-associated isozymes 65

Fenflumizole
HPLC assay of fenflumizole and its demethyl metabolites in biological samples 189

Flunoxaprofen
Determination of the non-steroidal anti-inflammatory drug flunoxaprofen, *S*(+)-2-(4-fluorophenyl)-α-methyl-5-benzoxazoleacetic acid, in blood and urine by HPLC 338

S(+)-2-(4-Fluorophenyl)-α-methyl-5-benzoxazoleacetic acid
Determination of the non-steroidal anti-inflammatory drug flunoxaprofen, *S*(+)-2-(4-fluorophenyl)-α-methyl-5-benzoxazoleacetic acid, in blood and urine by HPLC 338

Glycerol mononitrates
GC assay of glycerol mononitrates in biological samples 91

Glyceryl dinitrates
Determination of the two dinitrate metabolites of nitroglycerin in human plasma by capillary GC with electron-capture detection 101

Gramicidin S
Adsorption of cyclic peptides analogous to gramicidin S and gratisin onto octadecylsilica stationary phase and bacterial cells 287

Gratisin
Adsorption of cylic peptides analogous

to gramicidin S and gratisin onto octadecylsilica stationary phase and bacterial cells 287
Haemoglobins
Isolation and partial characterization of pyridoxal 5'-phosphate hemoglobins by HPLC as a quality-control method for hemoglobin-based blood substitutes 53
Histidine
Determination of stable isotopically substituted histidine in human plasma by GC–MS 1
Homogentisic acid
Analysis of homogentisic acid in body fluids by HPLC 233
Hyaluronic acid
Rapid and sensitive method for measurement of hyaluronic acid and isomeric chondroitin sulfates using HPLC 9
Hydrocarbons
Determination of hydroxylated metabolites of polycyclic aromatic hydrocarbons in urine 227
4-Hydroxyalprenolol
Derivatization of 2-amino alcohols with phosgene in aqueous media: limitations of the reaction selectivity as found in the presence of O-glucuronides of alprenol in urine 270
Hydroxyanisole
Sensitive HPLC method for the routine determination of butylated hydroxyanisole in plasma 282
2-Hydroxy-N-{3-[*m*-(1-piperidinylmethyl)-phenoxy]propyl}acetamide
Highly sensitive determination of 2-hydroxy-N-{3-[*m*-(1-piperidinylmethyl)-phenoxy]propyl}acetamide (desacetyl TZU-0460) in rat plasma by HPLC 370
p-Hydroxytriamterene
Rapid method for the determination of the diuretic triamterene and its metabolites in plasma and urine by HPLC 315
p-Hydroxytriamterene sulphate ester
Rapid method for the determination of the diuretic triamterene and its metabolites in plasma and urine by HPLC 315
Itraconazole
Determination of itraconazole in plasma and animal tissues by HPLC 332
Lipids
Method for determination of different phospholipids by circular HPTLC 257
Lipoperoxides
Determination of plasma lipoperoxides by HPLC 237
Lysophosphatidylcholine
Method for determination of different phospholipids by circular HPTLC 257
Methotrexate
Rapid and simple assay for the measurement of methotrexate in serum, urine and red blood cells be reversed-phase HPLC 320
3-Methylhistidine
Rapid HPLC of 3-methylhistidine in human urine 33
Nitroglycerin
Determination of the two dinitrate metabolites of nitroglycerin in human plasma by capillary GC with electron-capture detection 101
Oxymorphone
Electrochemical determination of oxymorphone in rat plasma by ion-pair reversed-phase HPLC 309
Oxytocin
Quantitative LC determination of disulfide-containing peptide analogous of vasopressin with dual Hg/Au electrochemical detection 43
Penicillins
Hospital routine analysis of penicillins, third-generation cephalosporins and aztreonam by conventional and high-speed HPLC 109
Peptides
Quantitative LC determination of disulfide-containing peptide analogous of vasopressin with dual Hg/Au electrochemical detection 43
Peptides, cyclic
Adsorption of cyclic peptides analogous to gramicidin S and gratisin onto octadecylsilica stationary phase and bacterial cells 287
Phosgene
Derivatization of 2-amino alcohols with phosgene in aqueous media: limitations of the reaction selectivity as found in the presence of O-glucuronides of alprenolol in urine 270
Phosphatidylcholine
Method for determination of different phospholipids by circular HPTLC 257
Phospholipids
Method for determination of different phospholipids by circular HPTLC 257

α-Phthalimidoglutarimide
Determination of thalidomide and its major metabolites by HPLC 181
Piperazine-2',5'-dione
LC determination of amoxicillin and its metabolites in human urine by post-column degradation with sodium hypochlorite 219
Polycyclic aromatic hydrocarbons
Determination of hydroxylated metabolites of polycyclic aromatic hydrocarbons in urine 227
Pyrazinamide
Rapid and simultaneous determination of pyrazinamide and its major metabolites in human plasma by HPLC 342
Pyridoxal 5'-phosphate haemoglobins
Isolation and partial characterization of pyridoxal 5'-phosphate hemoglobins by HPLC as a quality-control method for haemoglobin-based blood substitutes 53
Ranitidine
Determination of ranitidine in rat plasma and brain by HPLC 363
Saccharides
Rapid and sensitive method for measurement of hyaluronic acid and isomeric sulfates using HPLC 9
Sphingomyelin
Method for determination of different phospholipids by circular HPTLC 257
Sugar-chain heterogeneity
Sugar-chain heterogeneity of human alkaline phosphatases: differences between normal and tumour-associated isozymes 65
Sulindac
Analysis of sulindac and metabolites in plasma and urine by HPLC 171
Sulindac sulphide
Analysis of sulindac and metabolites in plasma and urine by HPLC 171
Sulindac sulphone
Analysis of sulindac and metabolites in plasma and urine by HPLC 171
Sulphate
Improved method for the determination of sulphate in human serum using ion chromatography 251
Tamoxifen
Quantitation of tamoxifen and N-desmethyltamoxifen in human plasma by HPLC, photochemical reaction and fluorescence detection, and its application to biopharmaceutic investigations 207
Thalidomide
Determination of thalidomide and its major metabolites by HPLC 181
Tocopherol
Determination of tocopherol and tocopherol acetate concentrations in human feces using HPLC 79
Tocopherol acetate
Determination of tocopherol and tocopherol acetate concentrations in human feces using HPLC 79
Triamterene
Rapid method for the determination of the diuretic triamterene and its metabolites in plasma and urine by HPLC 315
Trimethadione
Simultaneous determination of serum trimethadione and its metabolite by GC 376
Vasopressin
Quantitative LC determination of disulfide-containing peptide analogous of vasopressin with dual Hg/Au electrochemical detection 43
Vinpocetine
Quantitative determination of vinpocetine in human plasma by capillary GC–MS 264
Vitamin E
Determination of tocopherol and tocopherol acetate concentrations in human feces using HPLC 79
Vitamin H
Determination of free biotin in plasma by LC with fluorimetric detection 247